



## Fruit Tree Spraying Made Simple

### Fruits with Pits (Peaches, Apricots, Plums, Cherries, Nectarines)

To increase the effectiveness of all listed sprays, add *Spreader Sticker* at the rate of 1 Tablespoon per gallon to any mixed spray solution **except All Seasons Spray Oil**.

*Monterey Liqui-Cop* and *Bonide All Seasons Spray Oil* are natural sprays that are great for organic gardening.

WHEN	SPRAY	AMOUNT	INSECT/DISEASE
<b>January</b> When trees are dormant	<i>Bonide Liquid Copper</i>	<b>2 Tbsp/gal</b> except <b>4 Tbsp/gal</b> on cherries	Kills bacterial blights, leaf curl, and other fungal diseases. Spray the ground and branches thoroughly.
<b>February</b> When trees are dormant	<i>Bonide All Seasons Oil</i> (Do not mix with copper spray.)	<b>10 Tbsp/gal</b>	Kills overwintering insects (scale, mites, aphids, codling moth, whitefly, etc).
<b>March</b> When buds start to swell	<i>Bonide Liquid Copper @ 2 Tbsp/gal</i> OR <i>Bonide Fung-onil @ 1.5 Tbsp/gal</i>		Kills bacterial blights, leaf curl, and other fungal diseases. Spray the ground and branches thoroughly.
One week later	<i>Bonide Liquid Copper @ 2 Tbsp/gal</i> OR <i>Bonide Fung-onil @ 1.5 Tbsp/gal</i>		Kills bacterial blights, leaf curl, and other fungal diseases. Spray the ground, leaves and branches thoroughly.
After 2/3 of the flower petals have fallen	<i>Bonide Fruit Tree Spray</i>	<b>2.5 Tbsp/gal</b>	Keeps insects from attacking new fruit, prevents brown rot and blights.
<b>April</b>	<i>Bonide Fruit Tree Spray</i> (peaches, nectarines & cherries only)	<b>2.5 Tbsp/gal</b>	Keeps insects from attacking new fruit, prevents brown rot and blights.
<b>May</b>	<i>Bonide Fruit Tree Spray</i> (peaches, nectarines & cherries only)	<b>2.5 Tbsp/gal</b>	Keeps insects from attacking new fruit, prevents brown rot and blights.
<b>October or November</b> As soon as all the leaves fall off	<i>Bonide Liquid Copper @ 2 Tbsp/gal</i> except <b>4 Tbsp/gal</b> on cherries OR <i>Bonide Fung-onil @ 1 tsp/gal</i>		Kills bacterial blights, leaf curl, and other fungal diseases. Spray the ground and branches thoroughly.

Always consult and follow manufacturers' labels on all products.

1 oz = 2 Tbsp    1 Tbsp = 3 tsp

# Fruit Tree Spraying Made Simple

## Fruit with Seeds (Apples, Pears, Asian Pears)

To increase the effectiveness of all listed sprays, add *Bonide Turbo Spreader Sticker* at the rate of 1 Tablespoon per gallon to any mixed spray solution **except All Seasons Spray Oil**.

*Lime Sulfur* and *Bonide All Seasons Spray Oil* are natural sprays that are great for organic gardening.

### APPLES

WHEN	SPRAY	AMOUNT	INSECT/DISEASE
<b>January / February</b> When trees are dormant - no leaves or buds	<i>Lime Sulfur @ 8 Tbsp/gal combined</i> <i>All Seasons Spray Oil @ 2.5 Tbsp/gal</i> OR <i>Bonide All Seasons Spray Oil @</i> <i>10Tbsp/gal</i>		Kills scab and over-wintering insects.
<b>March</b> When branch tips show green tissue	<i>Bonide Fruit Tree Spray</i>	<b>1.5 Tbsp/gal</b>	Kills apple maggots, codling moth & other fruit damaging insects; rust, scab, mildew & assorted fruit rots. Spray the ground, branches and leaves thoroughly.
Two weeks later	<i>Bonide Fruit Tree Spray</i>	<b>2.5 Tbsp/gal</b>	Kills apple maggots, codling moth & other fruit damaging insects; aphids that cause distorted leaves; rust, scab, mildew & assorted fruit rots. Spray the ground, branches and leaves thoroughly.
<b>April, May or June</b> (depending on bloom time of specific variety)			
When most of the flower petals have fallen	<i>Bonide Fruit Tree Spray</i>	<b>2.5 Tbsp/gal</b>	Kills apple maggots, codling moth & other fruit damaging insects; rust, scab, mildew & assorted fruit rots. Spray the leaves and fruit thoroughly.
Two weeks after most of the flower petals have fallen	<i>Bonide Fruit Tree Spray</i>	<b>2.5 Tbsp/gal</b>	Kills apple maggots, codling moth & other fruit damaging insects; rust, scab, mildew & assorted fruit rots. Spray the leaves and fruit thoroughly.
Four weeks after most of the flower petals have fallen	<i>Bonide Fruit Tree Spray</i>	<b>2.5 Tbsp/gal</b>	Kills apple maggots, codling moth & other fruit damaging insects; rust, scab, mildew & assorted fruit rots. Spray the leaves and fruit thoroughly.

### PEARS and ASIAN PEARS

WHEN	SPRAY & AMOUNT	INSECT/DISEASE
<b>January / February</b> When trees are dormant - no leaves or buds	<i>Lime Sulfur @ 8 Tbsp/gal combined with</i> <i>All Seasons Spray Oil @ 2.5 Tbsp/gal</i> OR ONLY <i>Bonide All Seasons Spray Oil @ 7 Tbsp/gal</i>	Kills over-wintering insects.

Always consult and follow manufacturers' labels on all products.

1 oz = 2 Tbsp    1 Tbsp = 3 tsp

**541-673-8423**